

Description of G -bundles over G -spaces with quasi-free proper action of discrete group II

Morales Meléndez*, Quitzeh

December 27, 2009

1 The setting of the problem

This problem naturally arises from the Connor-Floyd's description of the bordisms with the action of a group G using the so-called fix-point construction. This construction reduces the problem of describing the bordisms to two simpler problems: a) description of the fixed-point set (or, more generally, the stationary point set), which happens to be a submanifold attached with the structure of its normal bundle and the action of the same group G , however, this action could have stationary points of lower rank; b) description of the bordisms of lower rank with an action of the group G . We assume that the group G is discrete.

Lets ξ be an G -equivariant vector bundle with base M .

$$\begin{array}{c} \xi \\ \downarrow \\ M \end{array}$$

Where the action of the group G is quasi-free over the base with normal stationary subgroup $H < G$ and there is no more fixed points of the action of the group H in the total space of the bundle ξ .

According [1, p.1] the bundle ξ separates as the sum of its G -subbundles:

$$\xi \approx \bigoplus_k \xi_k$$

where the index runs over all (unitary) irreducible representations $\rho_k : H \longrightarrow \mathbf{U}(V_k)$ of the group H and, as a H -bundle ξ_k can be presented as the tensor product:

$$\xi \approx \bigoplus_k \eta_k \otimes V_k,$$

*Partially supported by the grant 299388 of the mexican National Council for Science and Technology (CONACyT)

where the action of the group H over the bundles η_k is trivial, V_k denotes the trivial bundle with fiber V_k and with fiberwise action of the group H , defined using the linear representation ρ_k .

The particular case $\xi = \eta_k \otimes V_k$ was described in the previous article [1]. According [1, p.14] the bundle ξ_k can be obtained as the inverse image of a mapping

$$f_k : M/G_0 \longrightarrow B\text{Aut}_G(X_k)$$

where $G_0 = G/H$, $X_k = G_0 \times (F_k \otimes V_k)$ is the canonical model and $\text{Aut}_G(X_k)$ is the group of equivariant automorphisms of the space X_k as a vector G -bundle over the base G_0 . So, the bundle ξ can be given by a mapping

$$f : M/G_0 \longrightarrow \prod_k B\text{Aut}_G(X_k).$$

Consider the vector bundle over the discrete base G_0

$$X_\rho = G_0 \times \left(\bigoplus_k (F_k \otimes V_k) \right). \quad (1)$$

Define a fiberwise action $G \times X_\rho \rightarrow X_\rho$ by the formula

$$\begin{aligned} \phi([g], g_1) : [g] \times \left(\bigoplus_k (F_k \otimes V_k) \right) &\rightarrow [g_1 g] \times \left(\bigoplus_k (F_k \otimes V_k) \right) \\ \phi([g], g_1) &= \bigoplus_k (\text{Id} \otimes \rho_k(u(g_1 g)u^{-1}(g))) = \rho(u(g_1 g)u^{-1}(g)). \end{aligned} \quad (2)$$

Definition 1 *The bundle $X_\rho \rightarrow G_0$ with the just defined action is called the canonical model for the representation ρ .*

By $\text{Aut}_G(X_\rho)$ we denote the group of equivariant automorphisms of the canonical model X_ρ as a vector G -bundle over the base G_0 with fiber $\bigoplus_k (F_k \otimes V_k)$ and canonical action of the group G .

Lemma 1 *There exists a monomorphism*

$$i : \text{Aut}_G(X_\rho) \longrightarrow \prod_k \text{Aut}_G(X_k)$$

Proof. As before, an element of the group $\text{Aut}_G(X_\rho)$ is an equivariant mapping $\mathbf{A}^a$ such that the pair $(\mathbf{A}^a, a)$ defines a commutative diagram

$$\begin{array}{ccc} X_\rho & \xrightarrow{\mathbf{A}^a} & X_\rho \\ \downarrow & & \downarrow \\ G_0 & \xrightarrow{a} & G_0, \end{array}$$

$$a \in \text{Aut}_G(G_0) \approx G_0, \quad a[g] = [ga], \quad [g] \in G_0.$$

By the lemma 1 [1] applied to group of automorphisms $\text{Aut}_G(X_\rho)$, for $A^a \in \text{Aut}_G(X_\rho)$, we have

$$A^a|_{X_k} : X_k \longrightarrow X_k.$$

Note that this restriction is G -equivariant.

Define

$$i : \text{Aut}_G(X_\rho) \longrightarrow \prod_k \text{Aut}_G(X_k)$$

by the formula

$$i(A^a) = (A^a|_{X_k})_k.$$

This is clearly a homomorphism: it is a product of restrictions over invariant subspaces. Lets prove that it is injective. If $A^a|_{X_k} = \mathbf{Id}_{X_k}$, then $A^a = \mathbf{Id}_X$ because $X = \oplus_k X_k$.

In order to prove that the image of i is closed, note that it coincides with those automorphisms which commute with the inclusion

$$\Delta \times \mathbf{Id} : X_\rho \longrightarrow \prod_k X_k$$

i.e. if an element $(A_k^{a_k})_k \in \prod_k \text{Aut}_G(X_k)$ leaves the image of X_ρ invariant, then its restriction defines an element in $\text{Aut}_G(X_\rho)$ and the diagram

$$\begin{array}{ccc} X_\rho & \xrightarrow{A^a} & X_\rho \\ \downarrow & \searrow^{(A_k^{a_k})_k} & \downarrow \\ \prod_k X_k & \xrightarrow{(A_k^{a_k})_k} & \prod_k X_k \end{array}$$

commutes, i.e. $a^k = a \forall k$ and $A^a|_{X_k} = A_k^a$. In other words

$$\prod_k pr_k \circ i(A^a) = \Delta(a)$$

where

$$pr_k : \text{Aut}_G(X_k) \longrightarrow G_0$$

is the epimorphism of lemma 2 [1, p. 9] and

$$\Delta : G_0 \hookrightarrow \prod_k G_0.$$

Then

$$i(\text{Aut}_G(X_\rho)) = \prod_k pr_k^{-1} \Delta(G_0).$$

■

Corollary 1 *It takes place an exact sequence of groups*

$$1 \rightarrow \prod_k GL(F_k) \xrightarrow{\varphi} \text{Aut}_G(X_\rho) \xrightarrow{pr} G_0 \rightarrow 1$$

Proof. Define $pr = \Delta^{-1} \prod_k pr_k \circ i$. This is an epimorphism: let $A_k^a \in pr_k(a)$, then $(A_k^a)_k \in i(\text{Aut}_G(X_\rho))$.

We define the monomorphism

$$\varphi : \prod_k GL(F_k) \longrightarrow \text{Aut}_G(X_\rho)$$

using the monomorphisms

$$\varphi_k : GL(F_k) \longrightarrow \text{Aut}_G(X_k)$$

by the formula

$$\varphi = i^{-1} \prod_k \varphi_k.$$

Then $pr \circ \varphi = 1$ and if $a = 1$, then $\prod_k pr_k \circ i(A^1) = \Delta(1)$. This means that, for every k , there is a $B \in GL(F_k)$ such that $\varphi_k(B_k) = A^1|_{X^k}$, i.e. $\varphi(B_k)_k = A^1$. ■

Denote by $\text{Vect}_G(M, \rho)$ the category of G -equivariant vector bundles ξ over the base M with quasi-free action of the group G over the base and normal stationary subgroup $H < G$.

Then, by lemma 1 and the observations on p. 13 in [1] in terms of homotopy we have

$$\text{Vect}_G(M, \rho) \approx \prod_k [M, BU(F_k)] \quad (3)$$

Denote by $\text{Bundle}(X, L)$ the category of principal L -bundles over the base X .

Theorem 1 *There exists an inclusion*

$$\text{Vect}_G(M, \rho) \longrightarrow \text{Bundle}(M/G_0, \text{Aut}_G(X_\rho)). \quad (4)$$

Proof. We already have a monomorphism

$$\text{Vect}_G(M, \rho) \longrightarrow \text{Bundle}(M/G_0, \prod_k \text{Aut}_G(X_k))$$

i.e.

$$\prod_k \text{Vect}_G(M, \rho_k) \longrightarrow \prod_k \text{Bundle}(M/G_0, \text{Aut}_G(X_k))$$

see [1, p. 13]

A bundle $\xi \in \text{Vect}_G(M, \rho)$ is given by transition functions

$$\Psi_{\alpha\beta}(x) \in \prod_k \text{Aut}_G(X_k)$$

with the property that there exist $h_{\alpha,k}(x) \in G_0$ such that

$$h_{\alpha,k}^{-1}(x)pr_k \circ \Psi_{\alpha\beta}(x)h_{\alpha,k}(x)$$

does not depends on k . Lets show that can be found transition functions with the property that

$$\prod_k pr_k \Psi_{\alpha\beta}(x) = \Delta(a_{\alpha\beta}(x))$$

for some cocycle $a_{\alpha\beta}(x)$.

Because the group G_0 is discrete, for an atlas of connected charts with connected intersections, we can assume that $pr_k \circ \Psi_{\alpha\beta}(x) = a_{\alpha\beta,k}$ and $h_{\alpha,k}(x) = h_{\alpha,k} \in G_0$ do not depend on x and, therefore,

$$h_{\alpha,k}^{-1}(x)pr_k \circ \Psi_{\alpha\beta}(x)h_{\alpha,k}(x) = a_{\alpha\beta}$$

does not depend on k nor x . Let $H_{\alpha,k} \in \text{Aut}_G(X_k)$ such that $pr_k(H_{\alpha,k}) = h_{\alpha,k}$. Then,

$$\prod_k pr_k(H_{\alpha,k}^{-1} \Psi_{\alpha\beta}(x) H_{\alpha,k}) = \Delta(a_{\alpha\beta}).$$

■

Theorem 2 *If the space X is compact , then*

$$\text{Bundle}(X, \text{Aut}_G(X_\rho)) \approx \bigsqcup_{M \in \text{Bundle}(X, G_0)} \text{Vect}_G(M, \rho). \quad (5)$$

Proof. We will follow the proof theorem 3 in [1, p. 14]. Given a bundle $\xi \in \text{Bundle}(X, \text{Aut}_G(X_\rho))$ with transition functions

$$\Psi_{\alpha\beta}(x) \in \text{Aut}_G(X_\rho)$$

we obtain transition functions

$$pr \circ i \circ \Psi_{\alpha\beta}(x) \in \text{Aut}_G(G_0) \approx G_0,$$

defining an element $M \in \text{Bundle}(X, G_0)$ together with a projection $\xi \rightarrow M$. Changing the fibers $\text{Aut}_G(X_\rho)$ by X_ρ , we obtain an action of the group G , that reduces over the base to the factor group G_0 .

■

Lets rewrite this in terms of homotopy.

Corollary 2 *If the space X is compact, then*

$$[X, \text{Aut}_G(X_\rho)] \approx \bigsqcup_{M \in [X, BG_0]} \prod_k [M, BU(F_k)]. \quad (6)$$

■

2 The case when the subgroup $H < G$ is not normal

Consider an equivariant vector G -bundle ξ over the base M

$$\begin{array}{c} \xi \\ \downarrow p \\ M. \end{array}$$

Let $H < G$ be a finite subgroup. Assume that M is the set of fixed points of the conjugation class of this subgroup, more accurately

$$M = \bigcup_{[g] \in G/N(H)} M^{gHg^{-1}}, \quad (7)$$

and that there is no more fixed points of the conjugation class of H in the total space of the bundle ξ ; here we have denoted by M^H the set of fixed points of the action of the subgroup H over the space M , $N(H)$ the normalizer of the group H in G and we are using the equality $gM^H = M^{gHg^{-1}}$ and the fact that $lHl^{-1} = gHg^{-1}$ if and only if $g^{-1}l \in N(H)$.

Lets denote by $\mathfrak{F}_\xi$ the family of subgroups of G having non-trivial fixed points in the total space of the bundle ξ , i.e.

$$\mathfrak{F}_\xi = \{K < G \mid \xi^K \neq \emptyset\}.$$

This is a partial ordered set by inclusions and is closed under the action of the group G by conjugation¹. Also, the action

$$\begin{array}{ccc} G \times \mathfrak{F}_\xi & \longrightarrow & \mathfrak{F}_\xi \\ (g, K) & \mapsto & gKg^{-1} \end{array}$$

preserves the order.

Definition 2 *We will say that $H < G$ is the unique, up to conjugation, maximal subgroup for the G -bundle ξ if every conjugate gHg^{-1} is maximal in $\mathfrak{F}_\xi$ and there is no more maximal elements in this family.*

In this section will assume in any case, that $H < G$ is the unique, up to conjugation, maximal subgroup.

Lemma 2 *If $H \neq gHg^{-1}$, then*

$$M^H \cap M^{gHg^{-1}} = \emptyset$$

Proof. If there is an $x \in M^H \cap M^{gHg^{-1}}$ then, the point x is fixed under the action of the subgroup generated by H and gHg^{-1} , but this group is not contained in any of the subgroups of the form lHl^{-1} , $l \in G$. ■

¹If $\xi^K \neq \emptyset$, then $\xi^{gKg^{-1}} = g\xi^K \neq \emptyset$.

Lemma 3 *If the condition (7) holds, then the G -bundle ξ can be presented as a disjoint union of pair-wise isomorphic bundles with quasi-free action over the base. More precisely*

$$\xi = \bigsqcup_{[g] \in G/N(H)} \xi_{[g]},$$

where

$$\xi_{[g]} = p^{-1}(M^{gHg^{-1}})$$

is a vector bundle with quasi-free action of the group $N(gHg^{-1})$ and, for every element $g \in G$ the mapping

$$g : \xi^{[1]} \longrightarrow g\xi^{[1]} = \xi_{[g]}$$

defines an equivariant isomorphism of this bundles, i.e. the diagram

$$\begin{array}{ccc} N(H) \times \xi_{[1]} & \longrightarrow & \xi_{[1]} \\ \downarrow s_g \times g & & \downarrow g \\ N(gHg^{-1}) \times \xi_{[g]} & \longrightarrow & \xi_{[g]} \end{array} \quad (8)$$

commutes, where

$$s_g : N(H) \longrightarrow N(gHg^{-1}) = gN(H)g^{-1}, \quad (g, n) \mapsto gng^{-1}.$$

Proof. From lemma 2 it follows that

$$M = \bigsqcup_{[g] \in G/N(H)} M^{gHg^{-1}}$$

and, therefore,

$$\xi = \bigsqcup_{[g] \in G/N(H)} \xi_{[g]}.$$

Since the action of G is fiberwise, we have $g \cdot \xi_{[1]} = \xi_{[g]}$ for every $g \in G$. Restricting the projection $\xi \longrightarrow M$ to the space $\xi_{[g]}$, we obtain the bundle

$$\begin{array}{c} \xi_{[g]} \\ \downarrow p \\ M^{gHg^{-1}}. \end{array}$$

The bundle $\xi_{[g]}$ has an action of the normalizer $N(gHg^{-1})$:

$$N(gHg^{-1}) \times \xi_{[g]} \longrightarrow \xi_{[g]},$$

i.e. $\xi_{[g]}$ is a $N(gHg^{-1})$ -bundle for every $g \in G$.

Note that group conjugation $s_g : N(H) \longrightarrow N(gHg^{-1})$ defines an isomorphism between these groups that fits into the commutative diagram

$$\begin{array}{ccc} N(H) \times \xi_{[1]} & \longrightarrow & \xi_{[1]} \\ \downarrow & & \downarrow \\ N(gHg^{-1}) \times \xi_{[g]} & \longrightarrow & \xi_{[g]}. \end{array}$$

i.e. $gng^{-1} \cdot gx = g \cdot nx$. This means that the bundles $\xi_{[1]}$ and $\xi_{[g]}$ are naturally and equivariantly isomorphic.

Evidently, the mappings on the diagram (8) do not depend on the elements $n \in N(H)$, but they depend on the element $g \in G$.

The action of the group $N(H)$ over the base M^H reduces to the factor group $N(H)/H$:

$$\begin{array}{ccc} N(H) \times \xi_{[1]} & \longrightarrow & \xi_{[1]} \\ \downarrow & & \downarrow \\ N(H)/H \times M^H & \longrightarrow & M^H \end{array}$$

where, considering the maximality of the group H , the action $N(H)/H \times M \rightarrow M$ is free and, by hypothesis, there is no more fixed of the action of the subgroup H in the total space of the bundle ξ , i.e. $N(H)$ acts quasi-freely over the base and has normal stationary subgroup H . ■

Definition 3 *If the condition (7) holds, we will say that the group G acts quasi-freely over the bundle ξ with (non-normal) stationary subgroup H .*

As we will see in theorem 3, for classifying purposes, it is enough to consider bundles with normal stationary subgroup.

Let $X(\rho)$ be the canonical model for the representation $\rho : H \rightarrow GL(F)$ with action of the group $N(H)$. Define a canonical model $X(\rho_g)$ for the representation

$$\rho_g : gHg^{-1} \xrightarrow{s_g^{-1}} H \xrightarrow{\rho} GL(F),$$

$s_g(n) = gng^{-1}$. The action of the group $N(gHg^{-1})$ over $X(\rho_g)$ is defined using the homomorphism of right gHg^{-1} -modules

$$u_g : gHg^{-1} \xrightarrow{s_g^{-1}} H \xrightarrow{u} N(H) \xrightarrow{s_g} N(gHg^{-1})$$

by the formula (2).

Let

$$GX(\rho) := \bigsqcup_{[g] \in G/N(H)} X(\rho_g)$$

i.e. if $lHl^{-1} = gHg^{-1}$, then the spaces $X(\rho_g)$ and $X(\rho_l)$ coincide.

This notation will be clear after the next lemma.

Lemma 4 *The group G acts over the space $GX(\rho)$ quasi-freely with (non-normal) stationary subgroup H and, under this action, the space $GX(\rho)$ coincides with the orbit of the subspace $X(\rho)$. In particular, we have the relations*

$$N(H)(X(\rho)) = X(\rho)$$

and

$$(GX(\rho))^{gHg^{-1}} = N(gHg^{-1})/gHg^{-1}.$$

Proof. The action $G \times GX(\rho) \rightarrow GX(\rho)$ is defined in the following way: for a fixed $g \in G$ define the mapping

$$g : X(\rho) \longrightarrow X(\rho_g)$$

as

$$s_g \times \mathbf{Id} : N(H)_0 \times F \longrightarrow N(gHg^{-1})_0 \times F$$

($N(H)_0 = N(H)/H$) and, if $lHl^{-1} = gHg^{-1}$, then the mapping $l : X(\rho) \longrightarrow X(\rho_l)$ is chosen to make the diagram

$$\begin{array}{ccc} X(\rho_g) & \xrightarrow{s_g^{-1} \times \mathbf{Id}} & X(\rho) \\ \parallel & & \downarrow l^{-1}g \\ X(\rho_l) & \xrightarrow{l^{-1}} & X(\rho). \end{array} \quad (9)$$

commutative, i.e.

$$l = (s_g \times \mathbf{Id}) \circ (g^{-1}l)$$

where the mapping

$$g^{-1}l : X(\rho) \longrightarrow X(\rho) = X(\rho_{g^{-1}l}) \quad (10)$$

is the canonical left translation by the element $g^{-1}l \in N(H)$. ■

Corollary 3 *There is an isomorphism*

$$g : \text{Aut}_{N(H)}(X(\rho)) \xrightarrow{\approx} \text{Aut}_{N(gHg^{-1})}(X(\rho_g)) \quad (11)$$

that depends only on the class $[g] \in G/N(H)$.

Proof. We have a diagram (8) for $\xi = GX(\rho)$. Such a diagram always induces an isomorphism

$$\text{Aut}_{N(H)}(X(\rho)) \xrightarrow{\approx} \text{Aut}_{N(gHg^{-1})}(X(\rho_g))$$

by the rule

$$\mathbf{A} \mapsto g\mathbf{A}g^{-1}$$

and, if $l \in [g] \in G/N(H)$ then $l^{-1}g \in N(H)$ commutes with $\mathbf{A} \in \text{Aut}_{N(H)}(X(\rho))$. Therefore

$$g\mathbf{A}g^{-1} = g(g^{-1}l)(l^{-1}g)\mathbf{A}g^{-1} = g(g^{-1}l)\mathbf{A}(l^{-1}g)g^{-1} = l\mathbf{A}l^{-1}. \quad \text{■}$$

Definition 4 *The space $GX(\rho)$ is called the canonical model for the case when the subgroup $H < G$ is not normal.*

Lemma 5

$$\text{Aut}_G(GX(\rho)) \approx \text{Aut}_{N(H)}(X(\rho)) \quad (12)$$

Proof. By definition, an element of the group $\text{Aut}_G(X)$ is an equivariant mapping $\mathbf{A}^a$ such that the pair $(\mathbf{A}^a, a)$ defines the commutative diagram

$$\begin{array}{ccc} X & \xrightarrow{\mathbf{A}^a} & X \\ \downarrow & & \downarrow \\ G/H & \xrightarrow{a} & G/H, \end{array}$$

that commutes with the canonical action, i.e. the mapping $a \in \text{Aut}_G(G/H)$ satisfies the condition

$$a \in \text{Aut}_G(G/H) \approx N(H)/H, \quad a[g] = [ga], \quad [g] \in N(H)/H.$$

Therefore, $\mathbf{A}^a = (A^a[g])_{[g] \in N(H)/H} \in \text{Aut}_{N(H)}(X(\rho))$.

The value of the operators $(A^a[g])_{[g] \in G/H}$ can be calculated in terms of the operator $A^a[1]$ as in lemma 2 from [1, p. 9]. ■

Denote by $\widetilde{\text{Vect}}_G(M, \rho)$ the category of vector bundles with quasi-free action of the group G over the base M .

Theorem 3 $\widetilde{\text{Vect}}_G(M, \rho) \approx \text{Vect}_{N(H)}(M^H, \rho)$.

Proof. From lemma 3 follows that the bundles $\xi_{[1]}$ and $\xi_{[g]}$ equivariantly isomorphic and are given by mappings

$$M^{gHg^{-1}}/N(gHg^{-1})_0 \longrightarrow B\text{Aut}_{N(gHg^{-1})}(X(\rho_g)),$$

and

$$M^H/N(H)_0 \longrightarrow B\text{Aut}_{N(H)}(X(\rho)),$$

that can be put in the commutative diagram

$$\begin{array}{ccc} M^H/N(H)_0 & \longrightarrow & B\text{Aut}_{N(H)}(X(\rho)) \\ \downarrow \bar{g} & & \downarrow \bar{g} \\ M^{gHg^{-1}}/N(gHg^{-1})_0 & \longrightarrow & B\text{Aut}_{N(gHg^{-1})}(X(\rho_g)). \end{array}$$

Here, $g : \xi^H \longrightarrow \xi^{gHg^{-1}}$ is the action over the bundle ξ . The arrow on the right side is induced by the isomorphism (11) and does not depend on the element $g \in [g] \in G/N(H)$. ■

References

- [1] Mishchenko A.S., Morales Meléndez, Quitze. *Description of G-bundles over G-spaces with quasi-free proper action of discrete group*
- [2] Luke G., Mishchenko A. S., *Vector Bundles And Their Applications*. Kluwer Academic Publishers Group (Netherlands), 1998. ISBN: 9780792351542
- [3] P. Conner, E. Floyd. *Differentiable periodic maps*. Berlin, Springer-Verlag 1964.

- [4] Palais R.S. *On the Existence of Slices for Actions of Non-Compact Lie Groups* Ann. Math., 2nd Ser., Vol. 73, No. 2. (1961), pp. 295-323.
- [5] Atiyah M.F., *K-theory*. Benjamin, New York, (1967).
- [6] Serre J.P., *Représentations linéaires des groupes finis*. Hermann, Paris. 1967.
- [7] Levine M., Serpé C., *On a spectral sequence for equivariant K-theory* K-Theory (2008) 38 pp. 177–222
- [8] Beyl F.R., Tappe J., *Groups Extensions, Representations, and the Schur Multiplier*. Springer-Verlag (Berlin Heidelberg), 1982. ISBN 354011954X
- [9] Brown K.S. *Cohomology of groups*. Springer-Verlag (New York Heidelberg Berlin), 1982.
- [10] Eilenberg S., MacLane S. *Cohomology theory in abstract groups. II. Group extensions with a non-abelian kernel*. Ann. Math., 1947, (2) 48, p. 326—341.